arXiv:1212.3831v3 [cond-mat.mes-hall] 8 Feb 2013

Comment on "Reply to arXiv:1212.3831v1 'Comment on: Magnetotransport through graphene spin valves and its following works' " by Ding, Zhu, and Berakdar (arXiv:1302.0613v1)

Y. Zhou and M. W. Wu[*]

Hefei National Laboratory for Physical Sciences at Microscale and Department of Physics,
University of Science and Technology of China, Hefei, Anhui, 230026, China

(Dated: February 12, 2013)

In the reply by Ding *et al.*[1] to our previous comment,[2] they claimed that the formula we proposed "is a misconception resulting from adding an energy and momentum dependent function that has to vanish due to fundamental reasons" and the formula in their previous work[3–5] is "valid irrespective of whether the system is infinite or extended". In the following, we will demonstrate that their claim is wrong due to fundamental reasons.

They started the discussion from the formula of current given by Eq. (2) in Ref. 1 [see also Eq. (12.21) in Ref. 6],

$$I = \frac{ie}{2h}\int d\varepsilon\ \mathrm{Tr}\Big\{[f_L(\varepsilon)\hat{\Gamma}_L(\varepsilon) - f_R(\varepsilon)\hat{\Gamma}_R(\varepsilon)][\hat{G}^r(\varepsilon) - \hat{G}^a(\varepsilon)] + [\hat{\Gamma}_L(\varepsilon) - \hat{\Gamma}_R(\varepsilon)]\hat{G}^<(\varepsilon)\Big\}, \tag{1}$$

where all symbols with ˆ represent the matrices in the space formed by a complete basic set in the investigated system. For graphene, the basic set is usually chosen to be the one formed by the joint eigenstates of momentum $\mathbf{q}$ and pseudospin s (with $s = a, b$ corresponding to A and B sublattices, respectively). From the Dyson equation, they obtained

$$\hat{G}^r(\varepsilon) - \hat{G}^a(\varepsilon) = \hat{G}^r(\varepsilon)[\hat{g}^a(\varepsilon)^{-1} - \hat{g}^r(\varepsilon)^{-1}]\hat{G}^a(\varepsilon) + \hat{G}^r(\varepsilon)[\hat{\Sigma}^r(\varepsilon) - \hat{\Sigma}^a(\varepsilon)]\hat{G}^a(\varepsilon). \tag{2}$$

Based on the facts that $\hat{g}^a(\varepsilon)^{-1} - \hat{g}^r(\varepsilon)^{-1} = -2i\eta \to 0$ and $\hat{G}^{r,a}(\varepsilon)$ has neither isolated poles nor a branch cut if $\hat{\Sigma}^{r,a}(\varepsilon)$ is finite, they claimed that the first term in Eq. (2) cannot contribute to the current given by Eq. (1) no matter whether the system is finite or infinite.

Unfortunately, these authors overlooked an obvious fact that *all Green's functions and self-energies in the infinite system are matrices with infinite dimensions and hence the trace over the above term proportional to* η *can still be finite*. In fact, under the assumption adopted in all their works:[3–5] only atoms in the A sublattice are connected with the electrodes and the energy and momentum dependence of the self-energy can be neglected, i.e., $\Sigma^{r,a}_{\alpha,\mathbf{k}s,\mathbf{k}'s'}(\varepsilon) = \delta_{s,a}\delta_{s',a}\Sigma^{r,a}_{\alpha}/N$ with $\alpha = L, R$ for the left and right leads and N being the number of sites on A sublattice, one obtains

$$\begin{aligned}&\mathrm{Tr}\{\hat{\Gamma}_\alpha[\hat{G}^r(\varepsilon) - \hat{G}^a(\varepsilon)]\}\\ &= \Gamma_\alpha\mathcal{G}^r_a(\varepsilon)[\overline{g}^a_a(\varepsilon)^{-1} - \overline{g}^r_a(\varepsilon)^{-1}]\mathcal{G}^a_a(\varepsilon)\\ &\quad + \Gamma_\alpha\mathcal{G}^r_a(\varepsilon)(\Sigma^r - \Sigma^a)\mathcal{G}^a_a(\varepsilon),\end{aligned} \tag{3}$$

in which $\mathcal{G}^{r,a}_a(\varepsilon) = \frac{1}{N}\sum_{\mathbf{qq}'} G^{r,a}_{\mathbf{q}a,\mathbf{q}'a}(\varepsilon)$ and $\overline{g}^{r,a}_a(\varepsilon) = \frac{1}{N}\sum_{\mathbf{q}} g^{r,a}_{\mathbf{q}a,\mathbf{q}a}(\varepsilon)$. This means that the contribution to the current from the first term in Eq. (2) is proportional to the first term in Eq. (3) and hence the first term in Eq. (5) in Ref. 2, which has been demonstrated to be nonzero in the infinite system in our previous comment.[2] We would like to stress that the above claim is independent of the value of the self-energy, in contrast to the claim in the reply by Ding *et al.*.[1] Actually, the first term in Eq. (5) in Ref. 2 is just the spectral density of the system when $\hat{\Sigma}^{r,a} = 0$. However, it is logically incorrect in their reply to treat this fact as the evidence that the above term becomes zero when $\hat{\Sigma}^{r,a} \neq 0$.



FIG. 1: (Color online) $|\Gamma_0\overline{g}^r_a(\varepsilon)|$ against the energy ε for $D = 2.3t_g$ (the Debye cutoff) and $\Gamma^\uparrow_L = \Gamma^\downarrow_L = \Gamma_0 = 1.15t_g$.

We further demonstrate that the first term in Eq (2) even dominates the current and hence the tunnel magnetoresistance (TMR) for parameters used in the work by Ding *et al.*.[3–5] From Eq. (5) in Ref. 2, one obtains

$$\mathcal{G}^r_a(\varepsilon) - \mathcal{G}^a_a(\varepsilon) = \frac{\overline{g}^r_a(\varepsilon) - \overline{g}^a_a(\varepsilon)}{|1 - \Sigma^r\overline{g}^r_a(\varepsilon)|^2} + \frac{\overline{g}^r_a(\varepsilon)(\Sigma^r - \Sigma^a)\overline{g}^a_a(\varepsilon)}{|1 - \Sigma^r\overline{g}^r_a(\varepsilon)|^2}, \tag{4}$$

with $\Sigma^{r,a} = \pm i(\Gamma_L + \Gamma_R)/2$. In order to compare the relative importance of the two terms in the above equation, which just correspond to the two terms in Eq. (2), we plot

$|\Gamma_0 \overline{g}_a^r(\varepsilon)|$ in Fig. 1 in the case without spin polarization, i.e., $\Gamma_L^\uparrow = \Gamma_L^\downarrow = \Gamma_0$. Here we adopt the Debye prescription just as Ding *et al.*:[3–5] the cutoff energy is chosen to ensure the conservation of the total number of states in the Brillouin zone after linearization of the spectrum around the K point, which gives $D = 2.3t_g$. We also set $\Gamma_0 = 0.5D = 1.15t_g$, which is the largest self-energy used in their work.[3–5] Since this approach and in fact any approach with a cutoff are only valid in the energy regime far below the cutoff energy D, we only present the result with energy $|\varepsilon| < 0.4D$. From Fig. 1, it is seen that $|\Gamma_0 \overline{g}_a^r(\varepsilon)|$ is much smaller than 1 in most cases. This indicates that the first term in Eq. (4) and hence the first term in Eq. (2) indeed dominates the current and hence the TMR. Therefore, due to the erroneous omission of the above dominate term, the formula by Ding *et al.*[3–5] is incorrect in the extended infinite graphene they investigated and all their results are scientifically groundless. In addition, from Eq. (4), one also finds that $\mathcal{G}_a^r(\varepsilon) - \mathcal{G}_a^a(\varepsilon) \approx \overline{g}_a^r(\varepsilon) - \overline{g}_a^a(\varepsilon)$ in this case. This means the conductance given by Eq. (18) in Ref. 2 is almost proportional to the density of states of this system, which is a natural result in the weak coupling limit. Therefore, the linear dependence on the bias of the conductance from our formula, shown in Fig. 1 in Ref. 2, is an adding support of the correctness of our result.

Last but not least, in Fig. 2 in the reply,[1] Ding *et al.* presented the TMR from the approach using our suggestion in Ref. 2. From their figure, they claimed that the TMR from our approach becomes zero at zero bias. However, it is impossible to obtain such a result through our formula [Eq. (18) in Ref. 2]. This can be proved as follows. In the zero-bias limit, $|\Gamma_0 \overline{g}_a^r(\varepsilon)| = 0$, thus $\mathcal{G}_a^r(\varepsilon) - \mathcal{G}_a^a(\varepsilon) = \overline{g}_a^r(\varepsilon) - \overline{g}_a^a(\varepsilon)$ from Eq. (4). Then exploiting Eq. (18) in Ref. 2 and the definition of the TMR $[I(0) - I(\pi)]/I(0)$ with $I(0)$ $[I(\pi)]$ being the current in the parallel (antiparallel) configuration of the electrode magnetization, one obtains [here $\gamma_\sigma(\theta) = \Gamma_L^\sigma \Gamma_R^\sigma(\theta)/(\Gamma_L^\sigma + \Gamma_R^\sigma(\theta))$]

$$\mathrm{TMR} = \frac{\sum_\sigma [\gamma_\sigma(0) - \gamma_\sigma(\pi)]}{\sum_\sigma \gamma_\sigma(0)} = \frac{(\Gamma_L^\uparrow - \Gamma_L^\downarrow)^2}{(\Gamma_L^\uparrow + \Gamma_L^\downarrow)^2}. \tag{5}$$

Obviously, this TMR is nonzero for nonzero spin polarization ($\Gamma_L^\uparrow \neq \Gamma_L^\downarrow$) used in their reply.[1] Apparently these authors again made some numerical errors in their computation.

---

[*] Author to whom correspondence should be addressed; Electronic address: mwwu@ustc.edu.cn.

[1] K.-H. Ding, Z.-G. Zhu, and J. Berakdar, arXiv:1302.0613v1.

[2] Y. Zhou and M. W. Wu, arXiv:1212.3831.

[3] K.-H. Ding, Z.-G. Zhu, and J. Berakdar, Phys. Rev. B **79**, 045405 (2009).

[4] K.-H. Ding, Z.-G. Zhu, Z.-H. Zhang, and J. Berakdar, Phys. Rev. B **82**, 155143 (2010).

[5] K.-H. Ding, Z.-G. Zhu, and J. Berakdar, Phys. Rev. B **84**, 115433 (2011).

[6] H. Haug and A. -P. Jauho, *Quantum kinetics in Transport and Optics of Semiconductors* (Springer, Berlin, 1998).